

31 July 2008 MAA MathFest 1

Confounder Influence on Attributed Cases

Milo Schield
W. M. Keck Statistical Literacy Project
Augsburg College

MathFest 2008
 Madison, Wisconsin 31 July 2008
www.StatLit.org/pdf/2008SchieldMathFest6up.pdf

31 July 2008 MAA MathFest 2

Attributed Cases

Attributed cases are events that are attributable or attributed to an associated factor. These statistical events do not assert a causal relation.

- 438,000 deaths/year **due to** smoke/smoking.
- Deaths **attributed to** overweight/obese:
 2004: 400,000 deaths/year
 2007: 28,500 deaths/year

31 July 2008 MAA MathFest 3

Percentage of Cases that are Attributable

Suppose these are the death rates for two hospitals:

- 10% for the city research hospital
- 4% for the rural non-research hospital

Based on this data alone, one can say that

- “60% of the deaths at the city research hospital are attributable to that hospital.”

The math is simple:

- $\text{Excess} / \text{Exposed} = (10\% - 4\%) / 10\% = 60\%$

31 July 2008 MAA MathFest 4

Number of Cases that are Attributable

Suppose these are patients at two hospitals:

- 100,000 at the city research hospital
- 10,000 at the rural non-research hospital

The # of deaths (given 10% and 4% death rates):

- 10,000 deaths at the city research hospital
- 400 deaths at the rural non-research hospital.

6,000 (60%) of the 10,000 observed deaths at city hospital are attributable to the hospital.

31 July 2008 MAA MathFest 5

Decisions Based on Statistical Deaths

With this data, what should we do?



- Fire the managers at this “hospital of death”?
- Retrain the staff
- Recommend that patients avoid this hospital?
- Stop funding under-performing hospitals?
- Close under-performing hospitals?

31 July 2008 MAA MathFest 6

Association is not [always] Causation

Statistical associations can be due to various influences other than internal causation:

- Confounding – influence of associated factors.
- Assembly – the choice of definitions and groups.
- Randomness – most influential in small samples.
- Error/bias – subject, measurement or sampling bias.



31 July 2008 MAA MathFest 9

Percentage of Cases that are Attributable

Consider these confounder-adjusted death rates:

- 8% for the city research hospital
- 6% for the rural non-research hospital

Based on this data alone, one can say that

- "25% of the deaths at the city research hospital are attributable to that hospital."

The math is simple:

- $Excess / Exposed = (8\% - 6\%) / 8\% = 25\%$

31 July 2008 MAA MathFest 10

Number of Cases that are Attributable

Suppose these are patients at two hospitals:

- 100,000 at the city research hospital
- 10,000 at the rural non-research hospital

The # of deaths (given 8% and 6% death rates):

- 8,000 deaths at the city research hospital
- 600 deaths at the rural non-research hospital.

2,000 (25%) of the 8,000 deaths at city hospital are attributable to the hospital.

31 July 2008 MAA MathFest 11

of Cases Attributable: City Hospital Deaths

6K: 60% of 10K	Attributable to City Hospital <u>75% reduction</u>	2K	Explained by Confounder
4K: 40% of 10K		2K 25%	
	Common to both groups	6K: 75% of 8K	Allocation of 10K Deaths Before/After taking into account Patient Condition
Before		After	

31 July 2008 MAA MathFest 12

Conclusion

To help people analyze numbers in the news, Statistical Literacy must focus on:

- the influence of confounding on associations
- Specifically, the influence of confounding on events attributable to an associated factor.

Students are amazed that "attributed to" or "due to" is totally unrelated to any causal claim. Students are dismayed to learn that these numbers are so soft – so easily influenced by other factors.